



## King's Research Portal

DOI:

[10.1016/j.jacc.2017.12.040](https://doi.org/10.1016/j.jacc.2017.12.040)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Mayr, M., Gerszten, R., & Kiechl, S. (2018). Cardiovascular Risk Beyond Low-Density Lipoprotein Cholesterol. *Journal of the American College of Cardiology*, 71(6), 633-635. <https://doi.org/10.1016/j.jacc.2017.12.040>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Cardiovascular Risk Beyond Low-Density Lipoprotein Cholesterol

Manuel Mayr, MD, PhD,<sup>a</sup> Robert Gerszten, MD,<sup>b,c</sup> Stefan Kiechl, MD<sup>d</sup>

<sup>a</sup> King's British Heart Foundation Centre, King's College London, United Kingdom

<sup>b</sup> Division of Pulmonary, Critical Care and Sleep Medicine, Beth Israel Deaconess Medical Center, Boston, MA, USA

<sup>c</sup> Division of Cardiovascular Medicine, Beth Israel Deaconess Medical Center, Boston, MA, USA

<sup>d</sup> Department of Neurology, Medical University of Innsbruck, Austria

**Short title:** Cardiovascular Risk Beyond Low-Density Lipoprotein Cholesterol

**Word Count:** 2,031

**KEY WORDS** apolipoproteins, biomarkers, mass spectrometry, nuclear magnetic resonance spectroscopy, precision medicine

**Disclosures.** Prof. M. Mayr is a British Heart Foundation (BHF) Chair Holder (CH/16/3/32406) with BHF programme grant support (RG/16/14/32397). Dr. Gerszten is supported by NIH grants R01HL098280, U01DK048489, R01DK081572, U24DK112340 and DK 108159. This work was supported by an excellent initiative (Competence Centers for Excellent Technologies [COMET]) of the FFG (Austrian Research Promotion Agency): Research Center of Excellence in Vascular Ageing-Tyrol, VASCage (K-Project Nr. 843536) funded by BMVIT (Federal Ministry for Transport, Innovation and Technology), BMWFW (federal Ministry of Science, Research and Economy), the Wirtschaftsagentur Wien, and Standortagentur Tirol. MM and SK filed / licensed patents on cardiovascular biomarkers.

---

## Correspondence:

Prof. Manuel Mayr  
King's British Heart Foundation Centre  
King's College London  
125 Coldharbour Lane  
London SE5 9NU, UK  
Fax: +44 (0)20 7848 5298  
Telephone: +44 (0)20 7848 5446  
E-mail: [manuel.mayr@kcl.ac.uk](mailto:manuel.mayr@kcl.ac.uk)

---

Over the last 50 years, routine lipid parameters for risk prediction of cardiovascular disease (CVD) have not changed: clinical assays still rely on simple biochemical measurements of total triglycerides, total cholesterol and high-density lipoprotein cholesterol (HDL-C). Low-density lipoprotein cholesterol (LDL-C) is not always measured but calculated from non-HDL-C. Further, standard lipid tests quantify the cholesterol or triglyceride content of lipoproteins, without providing size-specific lipoprotein particle information. By contrast, nuclear magnetic resonance spectroscopy (NMR) provides a rapid method for distinguishing and quantifying a wide range of lipoprotein subclasses. Upon exposure to a magnetic field, distinct lipoprotein subclasses emit a unique signal that is directly proportional to their concentration. Although NMR lacks sensitivity when compared with mass spectrometry, it can be used to analyse the lipid composition of lipoprotein subclasses (1,2).

In this edition of the *Journal*, NMR-based findings from The China Kadoorie Biobank Study contribute to refining the quantitative and qualitative features of atherogenic lipid profiles (3). Very low-density lipoprotein (VLDL) particle concentrations were at least as strongly associated with myocardial infarction (MI) and ischaemic stroke (IS) as LDL particles. Further, triglycerides (TGs) were more consistently related with MI and IS across the entire spectrum of lipoprotein subfractions than cholesterol, and VLDL and remnant cholesterol outperformed LDL-C in CVD risk prediction in this Chinese population characterized with low mean LDL-C levels (85 mg/dL in the control group).

A causal role for remnant cholesterol in TG-rich lipoproteins such as VLDL has previously been suggested by meta-analysis and large-scale Mendelian Randomization studies (4,5). Similarly, mass spectrometry-based proteomics in the Bruneck Study ranked apoCIII, apoCII and apoE - three VLDL-associated apolipoproteins - as first to third with regards to CVD risk prediction (6). However, these data must be interpreted in the context of the widespread

use of statins for primary prevention in this cohort. Emerging studies suggest that cause for MI is shifting from plaque ruptures to plaque erosions, possibly due to the widespread use of LDL-C lowering therapies (7,8). With low LDL-C levels, the relative contribution of TG-rich lipoproteins to CVD risk may increase.

The China Kadoorie Biobank Study offers additional remarkable insights. First, the data contradict the conventional, yet outdated view that lipids are predominantly a risk factor for MI and are only a weak predictor for IS. In fact, the associations of lipoprotein particles with MI and IS were highly concordant, and their magnitude was only marginally lower for IS. This is supported by other studies that obtained similar findings for MI and IS, i.e. by measuring the plasma lipidome (9) or apolipoproteins like lipoprotein (a) (10-12). The most recent meta-analysis of LDL-C lowering therapies yielded a risk reduction of 24% for MI and 19% for overall stroke for more-intensive versus less-intensive therapy (13). The randomized controlled trials (RCTs) for PCSK9 inhibition and ezetimibe therapy but not cholesterylester transfer protein inhibition with anacetrapib reported benefits of lipid-lowering therapy for IS similar to or even higher than for MI (14-17). Taken together, these findings are consistent with the notion that lipids promote atherosclerosis systemically and that atherosclerosis is a main underlying cause for IS across its major subtypes. The pathogenetic relevance of lipids is obvious for IS as a downstream manifestation of atherosclerosis in extra- or intracranial large arteries, one of the main causes of stroke in China. It is also plausible for small vessel stroke. Small vessel strokes commonly arise from plaques at the orifice of the penetrating artery rather than from lipohyalinosis. The pathogenic relevance, however, may also extend to cardioembolic stroke and Embolic Stroke of Undetermined Source (ESUS). Stiffening of the aorta due to atherosclerosis and subsequent loss of the Windkessel function may elicit a diastolic backwards flow at the upper circumference of the aortic arch redirecting cardiac

emboli into the cerebral circulation. A large proportion of ESUS may actually be the consequence of fissuring of non-stenotic plaques that escape detection by carotid ultrasound. Further research into the role of lipids in the IS subtype is required to draw more definitive conclusions.

Second, the Chinese Kadoorie Biobank Study provides clarity as to one of the most controversial topics in stroke medicine - the purported protective role of lipids for intracerebral haemorrhage (ICH). In brief, none of 61 NMR parameters (lipoprotein particle concentrations and composition, particle size, and apolipoproteins) exhibited a significant relationship with ICH despite the high incidence of ICH in the Chinese population and the large number of cases (1138 ICH patients) included in the analysis. There was no signal suggesting that low cholesterol or triglyceride levels confer a higher risk of ICH as has been suggested by a large-scale meta-analysis of observational studies including 1.4 million participants and 7960 ICH cases (18). This literature-based meta-analysis may have been confounded by unrecorded comorbidities that alter lipid levels (e.g. liver and renal disease, inflammatory diseases and malignancies), inclusion of high-risk individuals for cardiovascular disease who were treated with statins, and other determinants of ICH risk (e.g. alcohol consumption and socio-economic status) that were not rigorously assessed. This view is corroborated by genetic association studies demonstrating a higher risk of ICH in carriers of variants related to high rather than low cholesterol levels (19). Moreover, the initial finding of the Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL) trial that high-dose statin therapy increases the risk for ICH appears to have been an artefact of the intention-to-treat principle and was not confirmed in any other RCT testing lipid-lowering strategies, nor in subsequent meta-analyses (20,21). Even in the setting of stroke thrombolysis, there was no higher risk of ICH among statin users (22). Although it is

premature to close this discussion, the study by Holmes and colleagues in this edition of the *Journal* adds further evidence that the proposed link between low cholesterol and ICH is a misconception.

Third, now that most RCTs on HDL-C raising strategies have failed and Mendelian Randomization studies refute the causal role of HDL-C in CVD, research should target HDL composition, functionality (e.g. cholesterol efflux capacity), and other qualitative features. The Chinese Kadoorie Biobank Study suggests that the TG rather than the cholesterol content of HDL is a determinant of atherogenicity. Finally, circulating glycoprotein N-acetyl glucosamine residues - a glycan biomarker linked to inflammation and aging - showed one of the strongest associations across all three main vascular endpoints (MI, IS, and ICH). More studies are required to scrutinize these associations and to further explore cause and effect.

We are entering a new era of lipid management (12, 14-16). With a growing armamentarium of lipid-lowering therapies, patients can be more readily treated to achieve the recommended LDL-C target levels. Besides LDL-C, the therapeutic focus may broaden to tackle the residual CVD risk and include VLDL and TGs (3-6,23), as well also fatty acid composition (9, 24) and other apolipoproteins, i.e. lipoprotein (a) (10-12). It is time to advance NMR and mass spectrometry technologies for lipoprotein and apolipoprotein profiling to meet the high-throughput, low cost and standardization required for potential clinical use (25, 26). The application of multi-omics technologies might pave the way towards redefining CVD risk (27).

## REFERENCES

- (1) Ussher JR, Elmariah S, Gerszten RE, Dyck JR. The emerging role of metabolomics in the diagnosis and prognosis of cardiovascular disease. *J Am Coll Cardiol*. 2016;68:2850-2870.
- (2) Hinterwirth H, Stegemann C, Mayr M. Lipidomics: quest for molecular lipid biomarkers in cardiovascular disease. *Circ Cardiovasc Genet*. 2014;7:941-54.
- (3) Holmes MV, Millwood IY, Kartsonaki C et al. Lipids, lipoproteins and metabolites and risk of incident myocardial infarction and stroke subtypes. *J Am Coll Cardiol* 2018; x:xxx-xx.
- (4) Sarwar N, Sandhu MS, Ricketts SL et al. Triglyceride-mediated pathways and coronary disease: collaborative analysis of 101 studies. *Lancet* 2010;375:1634-1639.
- (5) Nordestgaard BG. Triglyceride-rich lipoproteins and atherosclerotic cardiovascular disease: new insights from epidemiology, genetics, and biology. *Circ Res* 2016;118:547-563.
- (6) Pechlaner R, Tsimikas S, Yin X et al. Very-low-density lipoprotein-associated apolipoproteins predict cardiovascular events and are lowered by inhibition of APOC-III. *J Am Coll Cardiol* 2017;69:789-800.
- (7) Libby P, Pasterkamp G. Requiem for the 'vulnerable plaque'. *Eur Heart J* 2015;36:2984-2987.
- (8) Arbab-Zadeh A, Fuster V. The myth of the "vulnerable plaque": transitioning from a focus on individual lesions to atherosclerotic disease burden for coronary artery disease risk assessment. *J Am Coll Cardiol* 2015;65:846-855.
- (9) Stegemann C, Pechlaner R, Willeit P et al. Lipidomics profiling and risk of cardiovascular disease in the prospective population-based Bruneck study. *Circulation* 2014;129:1821-1831.
- (10) Erqou S, Kaptoge S, Perry PL et al. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA* 2009;302:412-423.
- (11) Willeit P, Kiechl S, Kronenberg F et al. Discrimination and net reclassification of cardiovascular risk with lipoprotein(a): prospective 15-year outcomes in the Bruneck Study. *J Am Coll Cardiol* 2014;64:851-860.
- (12) Viney NJ, van Capelleveen JC, Geary RS, et al. Antisense oligonucleotides targeting apolipoprotein(a) in people with raised lipoprotein(a): two randomised, double-blind, placebo-controlled, dose-ranging trials. *Lancet*. 2016;388(10057):2239-2253.
- (13) Koskinas KC, Siontis GCM, Piccolo R et al. Effect of statins and non-statin LDL-lowering medications on cardiovascular outcomes in secondary prevention: a meta-analysis of randomized trials. *Eur Heart J* 2017. Oct 23. doi: 10.1093/eurheartj/ehx566.

- (14) Cannon CP, Blazing MA, Giugliano RP et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;372:2387-2397.
- (15) Sabatine MS, Giugliano RP, Wiviott SD et al. Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med* 2015;372:1500-1509.
- (16) Ridker PM, Revkin J, Amarenco P et al. Cardiovascular efficacy and safety of Bococizumab in high-risk patients. *N Engl J Med* 2017;376:1527-1539.
- (17) Bowman L, Hopewell JC, Chen F et al. Effects of Anacetrapib in patients with atherosclerotic vascular disease. *N Engl J Med* 2017;377:1217-1227.
- (18) Wang X, Dong Y, Qi X, Huang C, Hou L. Cholesterol levels and risk of hemorrhagic stroke: a systematic review and meta-analysis. *Stroke* 2013;44:1833-1839.
- (19) Akoudad S, Ikram MA, Portegies ML et al. Genetic loci for serum lipid fractions and intracerebral hemorrhage. *Atherosclerosis* 2016;246:287-292.
- (20) Spence JD. Statins do not cause intracerebral hemorrhage. *Neurology* 2012;79:1076-1077.
- (21) Endres M, Nolte CH, Scheitz JF. Statin treatment in patients with Intracerebral Hemorrhage. *Stroke* 2017; Nov 30. pii: STROKEAHA.117.019322.
- (22) Scheitz JF, MacIsaac RL, Abdul-Rahim AH et al. Statins and risk of poststroke hemorrhagic complications. *Neurology* 2016;86:1590-1596.
- (23) Nordestgaard BG. A test in context: lipid profile, fasting versus nonfasting. *J Am Coll Cardiol.* 2017;70:1637-1646.
- (24) Rhee EP, Cheng S, Larson MG, et al. Lipid profiling identifies a triacylglycerol signature of insulin resistance and improves diabetes prediction in humans. *J Clin Invest.* 2011;121:1402-11.
- (25) Soininen P, Kangas AJ, Würtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ Cardiovasc Genet.* 2015;8:192-206.
- (26) Yin X, Baig F, Haudebourg E, Blankley RT, et al. Plasma proteomics for epidemiology: increasing throughput with standard-flow rates. *Circ Cardiovasc Genet.* 2017;10(6). pii: e001808.
- (27) Rader DJ, Bajaj A, Khetarpal SA. Deep apolipoprotein proteomics to uncover mechanisms of coronary disease risk. *J Am Coll Cardiol.* 2017;69:801-804.